

Budislav Tatić & Zoran Krivošej

### ***Tulipa serbica (Liliaceae), a new species from Serbia***

#### **Abstract**

Tatić, B. & Krivošej, Z.: *Tulipa serbica (Liliaceae)*, a new species from Serbia. – Bocconeia 5: 733-736. 1997. – ISSN 1120-4060.

A new species of *Tulipa*, related to and previously confused with *T. scardica*, is described from Serbia (Yugoslavia). It is confined to serpentine soils and flowers in April to May.

According to the literature (Pančić 1884, Diklić 1975), two wild species of *Tulipa* grow on the territory of Serbia, *T. hungarica* Borb. (*T. billietiana* sensu Pančić) and *T. sylvestris* L. In addition, Pavlović (1962) recorded *T. scardica* Bornm. from Mt Rogozna. We found some specimens of *Tulipa* on the hill Beli Laz near the village Srbovac, a locality situated near Mt Rogozna but on the opposite side of the river Ibar.

We also visited the locus classicus of *Tulipa scardica*, on hills near the village Raduše in the F.Y.R. Makedonija, where we collected specimens of that species. We found differences between *T. scardica* (see Bornmüller 1923, Mayer & Micevski 1970) and our plants from Beli Laz as well as those from Mt Rogozna that Pavlović (1962) had accepted to be Bornmüller's species. In an earlier note (Tatić & al. 1991) we already drew attention to these differences. We have since then made a detailed study of *T. scardica*, based on literature data, herbarium specimens (BEO, W, SOM), and live material. We conclude that our Serbian specimens belong to a species new to science, described and named here.

***Tulipa serbica* Tatić & Krivošej, sp. nova.** – Holotypus in herbario Prirodno-Matematicki Fakultet, Priština, adservatur. – Fig. 1.

Bulbus ovatus, tunicis brunneis intus appresse pilosus; caulis 10-40 cm altus, glaber, striatus, uniflorus; folia 3-5, pallide viridia, inferiora 10-30 cm longa et 1.5-3 cm lata plerumque undulata, superiora anguste linearia; perigonum campanulatum vel infundibuliforme-campanulatum, 2-5 cm longum, tepalis utrinque purpureis vel pallide vinaceo-purpureis, ad marginem albescensibus, intus macula carentibus, externis acuminatis, internis apicem versus crenulatis; stamina tepalis duplo breviora, filamento glabro pallido vel flavido, anthera violacea apice acuta; capsula obovata, loculicida, 2-3 cm longa et 1.5-2 cm lata. Solo serpentinico. – Habitat in collibus serpentinicis Serbie australis (Fig. 2),



Fig. 1. *Tulipa serbica*, general habit and details. – Drawn from a plant of the type collection.




Fig. 2. Map of Serbia, showing the total known distribution of *Tulipa serbica*. — ■ Mt Rogozna;  
● hill Beli Laz.

ad 500-950 m supra maris aequor, ubi floret mensibus aprilie et maij. – Etymoogia: species patriae nostrae reipublicae Serbiae dicata.

*Specimens seen.* – Serbia, Mt Rogozna near Donja Kamenica, 900 m, on serpentine soil (herb. Priština; holotype); hill Beli Laz near Srbovac, 5 km N. of Kosovska Mitrovica, 900 m, on serpentine soil (herb. Priština).

*Tulipa serbica* is closely related to *T. scardica*, but differs in its paler, unspotted perianth segments, pale (not blackish) staminal filaments, and dull violet (not yellowish), acute anthers.

#### Acknowledgement

We would like to thank Dr S. Kožuharov for his help.

#### References

- Bornmüller, J. 1923: Über eine neue Tulpe der Flora Mazedoniens. – Repert. Spec. Nov. Regni Veg. **19**: 199-200.  
Diklić, N. 1975: 14 rod *Tulipa* L. – Pp. 534-537 in: Josifović, M. (ed.), Flora SR Srbije, **7**. – Beograd.  
Mayer, E. & Micevski, K. 1970: Zur Taxonomie und Chorologie von *Tulipa scardica* Bornm. – Feddes. Repert. **80**: 591-598.  
Pavlović, Z. 1962: Karakteristični elementi serpentinske flore Srbije. – Glasn. Prir. Muz. u Beogradu, Ser. B, Biol. Nauke **18**: 3-20.  
Pančić, J. 1884: Dodatak "Flori Kneževine Srbije". – Beograd.  
Tatić, B., Ružić, M. & Krivošej, Z. 1991: *Tulipa scardica* Bornm. nova vrsta u flori Srbije – Pp. 61-64 in: Anonymous (ed.), Zbornik radova sa Simpozijuma "Nedeljko Košanin i biološke nauke". – Ivanjica.

#### Addresses of the authors:

- Dr B. Tatić, Botanical Institute and Garden, Faculty of Science, University of Belgrade, Studentski Trg 16, YU-11000 Beograd, Yugoslavia.  
Mr Z. Krivošej, Faculty of Science, Department of Biology, Vidovdanska, JU-38000 Pristina, Yugoslavia.